

Stochastic Evolution of Graphs Using Local Moves

Hal Finkel

Perimeter Institute for Theoretical Physics

Department of Physics
Drexel University

LOOPS '05



Motivation

Questions

Definition of Local Moves

Outputs

Measurements

Conclusions




Questions

- ▶ Does a manifold-like structure result from the local application of Pachner moves?
- ▶ More generally, what are the properties of graphs generated by the local application of Pachner moves?




Expansion Move

An expansion move transforms a single vertex into a d -simplex (shown here for $d = 3$).



Contraction Move

A contraction move transforms a d -simplex into a single vertex (shown here for $d = 3$).



Exchange Move

An exchange move swaps the endpoints of two edges adjacent to some chosen edge.



Exchange Move and Embedding

Generalized exchange moves do not preserve planarity or any manifold-like structure.



Non-Local Edges

An edge is said to be “non-local” if it is not part of any short cycle (where short means the length of the cycle is $\leq r_l$). Intuitively, this should mean the edge connects locations in the graph which are “far away”.



Simulation Input Parameters

- ▶ d : The number of vertices in the simplex added during an expansion move.
- ▶ r_l : The length of the largest cycle to consider local.
- ▶ R : The ratio of exchange moves to expansion moves.
- ▶ R_c : The ratio of contraction moves to expansion moves.



Graph Pictures Key

The following graph pictures were drawn using an algorithm which treats each edge as a spring with some spring constant and solves for the equilibrium configuration.


The edges are colored as follows (where $r_l = 5$):

- ▶ Red - Initial edges which are “local”
- ▶ Magenta - Initial edges which are “non-local”
- ▶ Black - Edges added by expansion moves which are “local”
- ▶ Green - Edges added by expansion moves which are “non-local”




Initial Conditions

Initial conditions which were tried included a single triangle, regular lattices, almost complete graphs and random connected graphs (shown here with $n = 200$ and $m = 400$).




$$R = 0$$

With $R = 0$, the random graph has evolved to where $n = 2200$ and $m = 3400$. 14.7% of the edges are “non-local”.



$R = 100$

With $R = 100$, the random graph has evolved to where $n = 1800$ and $m = 2800$. 42.5% of the edges are “non-local”.



Statistical Dimension

Let $V(r, v)$ be the number of vertices within a distance r of a vertex v . If $V(r, v) = c_v r^{d_v}$ where c_v and d_v are some constants then d_v is the statistical dimension with respect to the vertex v . The statistical dimension for the entire graph, d , is obtained by averaging d_v for all vertices.



Analytic Formula

Average vertex degree:

$$\bar{d} = \frac{2m}{n}$$

Number of vertices within a distance r (“volume”):

$$V(r) = V(r-1) + A(r)$$

Number of vertices not within a distance r :

$$R(r) = n - V(r-1)$$



Analytic Formula (cont.)

Number of “free” edges at a distance r (which connect vertices at a distance r to those at a distance $r + 1$):

$$f(r) = (\bar{d} * A(r) - f(r-1)) \left(1 - \frac{1}{2} \left(\frac{A(r-1) - 1}{R(r) - 1} \right)\right)$$

Average number of “free” edges per vertex at a distance $r - 1$:

$$s(r) = f(r-1)/A(r-1)$$


The number of vertices at a distance r :

$$A(r) = R(r) \left(1 - \left(1 - \frac{s(r)}{R(r)}\right)^{A(r-1)}\right)$$




Analytic Formula Results

Plot of $\ln(V(r))$ vs. $\ln(r)$ for random connected graphs:




$R = 1$ Statistical Dimension Information

Plot of $\ln(V(r))$ vs. $\ln(r)$ for evolving graph with $R = 1$:




$R = 100$ Statistical Dimension Information

Plot of $\ln(V(r))$ vs. $\ln(r)$ for evolving graph with $R = 100$:



Small-Scale Dimension

Plot of small-scale dimension vs. R:




Small-Scale Dimension (cont.)

It seems that after some value of R the dynamics becomes “exchange dominated” whereby further increasing R does not significantly change the structure of the graph. This can also be seen using other measures (such as graph diameter).




Asymptotic Fraction of Non-Local Edges

A plot of the fraction of non-local edges vs. n :




Effect of r_l

For a graph evolved to a fixed size $n = 5400$, a plot of the fraction of non-local edges vs. r_l :



Effect of r_l (cont.)

For a graph evolving at $R = 100$, a plot of the fraction of non-local edges vs. r_l :



Conclusions

- ▶ Stochastic application of local moves produces graphs which do not seem directly interpretable as approximating a smooth manifold. Thus, if graphs of similar structure represent space then it will be necessary to use some dynamic probe to characterize the approximated manifold.
- ▶ Unlabeled graphs will not suffice, additional degrees of freedom seem to be necessary in order to construct local moves which produce more manifold-like graphs.
- ▶ Macroscopic locality may be fundamentally non-local.



Acknowledgments

- ▶ LOOPS '05 Organizing Committee
- ▶ Perimeter Institute for Theoretical Physics
 - ▶ Lee Smolin
 - ▶ Fotini Markopoulou-Kalamara
 - ▶ Maya Paczuski
 - ▶ Peter Grassberger
- ▶ Drexel University

